

Modularity in orbifold theory for vertex operator superalgebras

Chongying Dong¹ and Zhongping Zhao

Department of Mathematics, University of California, Santa Cruz, CA 95064

Abstract

This paper is about the orbifold theory for vertex operator superalgebras. Given a vertex operator superalgebra V and a finite automorphism group G of V , we show that the trace functions associated to the twisted sectors are holomorphic in the upper half plane for any commuting pairs in G under the C_2 -cofinite condition. We also establish that these functions afford a representation of the full modular group if V is C_2 -cofinite and g -rational for any $g \in G$.

1 Introduction

Modular invariance of trace functions is fundamental in the theory of vertex operator algebras and conformal field theory (see for example [B2], [DVVV], [DLM3], [FLM3], [H], [Hu], [M2], [MS], [V], [Z]). Many important results such as the existence of twisted sectors in orbifold theory, the Lie algebra structure of weight one subspace of a good vertex operator algebra, and classification of holomorphic vertex operator algebras with small central charges can be obtained by studying the modular invariance property of 1-point functions on torus (see [DM1], [DM2], [Mo], [S]). The modular invariance of trace functions associated to modules for an arbitrary rational vertex operator algebra was first established under the C_2 -cofinite condition in [Z]. This work was extended in [DLM3] to include a finite automorphism group of the vertex operator algebra. There were further generalizations in [M1], [M2], [Y].

Partially motivated by the generalized moonshine conjecture [N], the trace functions associated to twisted modules for a vertex operator algebra V and a finite automorphism group G was studied in [DLM3]. Under some finiteness conditions on a rational vertex operator algebra V , among other things, the precise number of inequivalent, simple g -twisted V -modules was determined and the modular invariance (in a suitable sense) of the 1-point functions on torus associated to any commuting pairs in G was established. If V is holomorphic, the existence and the uniqueness of the twisted sector for each $g \in G$ was obtained. The important case in which $V = V^\natural$ is the moonshine vertex operator algebra [FLM3] and G is the monster simple group [G] plays a special role in a proof of several parts of the generalized moonshine conjecture [DLM3] except the genus zero property. Also see [T1], [T2] and [T3] on the generalized moonshine conjecture.

¹Supported by NSF grants, a China NSF grant and a Faculty research grant from the University of California at Santa Cruz.

In this paper we present a general theory of modular invariance of trace functions associated to twisted sectors for a vertex operator superalgebra V and a finite automorphism group G . Note that V has a canonical central automorphism σ arising from the superspace structure of V , which is crucial in formulating the main results in this paper. Let $\bar{G}$ be the group generated by G and σ . We now explain the main results in details.

Let $g \in G$ have order T and σg have order T' . Then a simple σg -twisted module $M = (M, Y_M)$ has a grading of the shape

$$M = \bigoplus_{n=0}^{\infty} M_{\lambda+n/T} \quad (1.1)$$

with $M_\lambda \neq 0$ for some complex number λ which is called the *conformal weight* of M (see Section 4 and [DZ]). One basic result in [DZ] is that if V is g -rational there are only finitely many simple σg -twisted V -modules up to isomorphism. As in [DLM3], for any $h \in \bar{G}$ we can define a $h\sigma gh^{-1}$ -twisted module $h \circ M$ so that $h \circ M = M$ as vector spaces and $Y_{h \circ M}(v, z) = Y_M(h^{-1}v, z)$ (see Section 6). M is called h -stable if M and $h \circ M$ are isomorphic. As a result the stabilizer of M in $\bar{G}$ acts projectively on M , denoting by ϕ .

Let $v \in V$ and M be σh -stable. We set

$$T_M(v, (g, h), q) = q^{\lambda-c/24} \sum_{n=0}^{\infty} \text{tr}_{M_{\lambda+n/T}}(o(v)\phi(\sigma h))q^{n/T} \quad (1.2)$$

where $Y_M(v, z) = \sum_{m \in \frac{1}{T}\mathbb{Z}} v(m)z^{-m-1}$ and $o(v) = v(\text{wt}v - 1)$ induces a linear map on each homogeneous subspace of M . We extend the notation $T_M(v, (g, h), q)$ to all $v \in V$ linearly.

As in [Z], there is another vertex operator superalgebra structure on V (see Section 3). If the original vertex operator superalgebra is defined over sphere, the second vertex operator superalgebra can be regarded as defined over a torus. We will denote the corresponding weight by $\text{wt}[v]$. Here are our main results on the modular invariance.

Theorem 1 *Suppose that V is a C_2 -cofinite vertex operator superalgebra and G a finite group of automorphisms of V .*

(i) *Let $g, h \in G$ and M be a simple, σh -stable, σg -twisted V -module. Then the trace function $T_M(v, (g, h), q)$ converges to a holomorphic function in the upper half plane $\mathfrak{h}$ where $q = e^{2\pi i\tau}$ and $\tau \in \mathfrak{h}$.*

(ii) *Suppose in addition that V is x -rational for each $x \in \bar{G}$. Let $v \in V$ satisfy $\text{wt}[v] = k$. Then the space of (holomorphic) functions in $\mathfrak{h}$ spanned by the trace functions $T_M(v, (g, h), \tau)$ for all choices of g, h in G and σh -stable M is a (finite-dimensional) $SL(2, \mathbb{Z})$ -module such that*

$$T_M|\gamma(v, (g, h), \tau) = (c\tau + d)^{-k} T_M(v, (g, h), \gamma\tau),$$

where $\gamma \in SL(2, \mathbb{Z})$ acts on $\mathfrak{h}$ as usual.

More precisely, if $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ then we have an equality

$$T_M(v, (g, h), \frac{a\tau + b}{c\tau + d}) = (c\tau + d)^k \sum_W \gamma_{M,W} T_W(v, (g^a h^c, g^b h^d), \tau),$$

where W ranges over the $\sigma g^a h^c$ -twisted sectors which are $g^b h^d$ and σ -stable. The constants $\gamma_{M,W}$ depend only on M, W and γ only.

Theorem 2 *Let V be a C_2 -cofinite and σ -rational vertex operator superalgebra. Let $\{M^1, \dots, M^n\}$ be the inequivalent σ -twisted V -modules which are σ -stable. Then for any $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in $SL(2, \mathbb{Z})$ there are constants γ_{ij} for $1 \leq i, j \leq n$ such that for any $v \in V$ with $\text{wt}[v] = k$,*

$$T_{M^i}(v, (1, 1), \frac{a\tau + b}{c\tau + d}) = (c\tau + d)^k \sum_{j=1}^n \gamma_{ij} T_{M^j}(v, (1, 1), \tau).$$

It is important to point out that Theorem 2 is an analogue of the main theorem in [Z]. In fact, if $V_1 = 0$ then V is a vertex operator algebra and $\sigma = id_V$. In this case, Theorem 2 reduces the main theorem in [Z]. In particular, the space of characters of irreducible V -modules is invariant under the action of the modular group. But if $V_1 \neq 0$ the space of characters of irreducible V -modules is no longer invariant under the action of the modular group. However, the space of σ -trace on σ -twisted sectors is modular invariant. So the automorphism σ plays a fundamental role in the study of modular invariance for vertex operator superalgebra and its orbifold theory. This will be illustrated in Section 9 with examples.

There is an assumption in both Theorems 1 and 2 on twisted modules. That is, the twisted modules are required to be σ -stable. There are examples in which irreducible σ -twisted modules are not σ -stable. So these modules which are not σ -stable are excluded in the discussion of modular invariance. To include these modules, one can use the Theta group $\Gamma_\theta = \Gamma(2) \cup \Gamma(2)S$ instead of the full modular group $SL(2, \mathbb{Z})$. Here $\Gamma(2)$ is the congruence subgroup of $SL(2, \mathbb{Z})$ consisting of elements which are congruent to the identity modulo 2 and $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. In fact if $G = 1$ this has been done in [H]. The σ does not play any role in this approach.

Again Let G be a finite automorphism group of V and $g \in G$. Let M be a simple g -twisted V -module. Let G_M be the stablizer of M in G . Then M is h -stable for any $h \in G_M$ and G_M acts projectively on M . As before we use ϕ to denote the projective representation. For $h \in G_M$ and $v \in V$ set

$$F_M(v, (g, h), q) = \text{tr}_M o(v) \phi(h) q^{L(0)-c/24}.$$

The analogue of Theorem 1 is the following which extends the modular invariance result in [H] to the orbifold theory.

Theorem 3 *Suppose that V is a C_2 -cofinite vertex operator superalgebra and G a finite group of automorphisms of V .*

(i) Let $g, h \in G$ and M be a simple, h -stable g -twisted V -module. Then the trace function $F_M(v, (g, h), q)$ converges to a holomorphic function in the upper half plane $\mathfrak{h}$.

(ii) Suppose in addition that V is x -rational for each $x \in G$. Let $v \in V$ satisfy $\text{wt}[v] = k$. Then the space of (holomorphic) functions in $\mathfrak{h}$ spanned by the trace functions $F_M(v, (g, h), \tau)$ for all choices of g, h in G and h -stable M is a (finite-dimensional) Γ_θ -module. That is, if $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_\theta$ then we have an equality

$$F_M(v, (g, h), \frac{a\tau + b}{c\tau + d}) = (c\tau + d)^k \sum_W \gamma_{M,W} F_W(v, (g^a h^c, g^b h^d), \tau),$$

where W ranges over the $g^a h^c$ -twisted sectors which are $g^b h^d$ -stable. The constants $\gamma_{M,W}$ depend only on M, W and γ only.

This paper is a super version of the theories developed in [DLM3]. The main ideas and the broad outline of our proof follow from those in [DLM3], [Z] and [H]. Two important results on g -rationality and the associative algebras $A_g(V)$ in [DZ] (also see Section 4) also play basic roles in this paper. If V is g -rational, then $A_g(V)$ is semisimple and the category of V -modules and the category of finite dimensional $A_g(V)$ -modules are equivalent. Furthermore there are finitely many irreducible g -twisted modules. These results enable us to prove that the space of 1-point (g, h) functions is finite dimensional. In fact, from the proof of the main results in this paper one can see that we only need the assumptions that V is C_2 -cofinite and $A_g(V)$ is a semisimple associative algebra. Although there are a lot of expectations on the relation among rationality, C_2 -cofiniteness, the semisimplicity of $A_g(V)$, we cannot remove either the C_2 -cofinite condition or the $A_g(V)$ semisimplicity condition.

This paper is organized as follows. In Section 2 we review several family of modular functions and elliptic functions, and their transformation laws following [DLM3]. Section 3 is about the definition of vertex operator superalgebra and vertex operator superalgebra on torus. In Section 3 we define weak, admissible and ordinary g -twisted module for a vertex operator superalgebra V and a finite order automorphism g . We also present the main results concerning the associative algebra $A_g(V)$ and g -rationality. We define the space of abstract 1-point (g, h) -functions in Section 5 and establish that the elements in the full modular group transform the space of 1-point (g, h) functions to another space of 1-point functions. We also discuss the shapes of q -expansions of the 1-point functions and prove that these functions satisfy differential equations with regular singularity. This leads us to the definition of formal 1-point (g, h) functions. Sections 6 and 8 are the key sections in this paper. We show in Section 6 that the σh -trace on σg -twisted modules produce 1-point (g, h) -functions. In fact, the inequivalent simple σg -twisted modules give linearly independent 1 point functions. In Section 8 we show that if V is σg -rational then the space of 1-point (g, h) functions has a basis consisting of the σh -trace on the inequivalent simple σg -twisted modules. We also give several important corollaries in Section 8 concerning the rationality of the central charges and the conformal weights of simple g -twisted modules. In Section 9 we present the modular invariance result on Γ_θ . We give an existence result on the twisted sectors in Section 7 and discuss several examples in Section 10 to illustrate the main theorems.

2 P -functions and Q -functions

In this section, we review some properties of the P -functions and Q -functions following [DLM3]. These functions will be used extensively in later sections.

Recall that the *modular group* $\Gamma = SL(2, \mathbb{Z})$ acts on the upper half plane $\mathfrak{h} = \{z \in \mathbb{C} | \text{Im} z > 0\}$ via Möbius transformations

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \tau = \frac{a\tau + b}{c\tau + d}. \quad (2.1)$$

Γ also acts on the right of $S^1 \times S^1$ via

$$(\mu, \lambda) \begin{pmatrix} a & b \\ c & d \end{pmatrix} = (\mu^a \lambda^c, \mu^b \lambda^d). \quad (2.2)$$

Let $\mu = e^{2\pi i j/M}$ and $\lambda = e^{2\pi i l/N}$ for integers j, l, M, N with $M, N > 0$. For each integer $k = 1, 2, \dots$ and each (μ, λ) we define a function P_k on $\mathbb{C} \times \mathfrak{h}$ as follows

$$P_k(\mu, \lambda, z, q_\tau) = P_k(\mu, \lambda, z, \tau) = \frac{1}{(k-1)!} \sum'_{n \in \frac{j}{M} + \mathbb{Z}} \frac{n^{k-1} q_z^n}{1 - \lambda q_\tau^n} \quad (2.3)$$

In the above equation, $q_x = e^{2\pi i x}$ and $\sum'$ means omit the term $n = 0$ when $(\mu, \lambda) = (1, 1)$. Then P_k converges uniformly and absolutely on compact subsets of the region $|q_\tau| < |q_z| < 1$. We have the following transformation properties (see Theorem 4.2 of [DLM3]).

Theorem 2.1 *Suppose that $(\mu, \lambda) \neq (1, 1)$. Then the following equalities hold:*

$$P_k(\mu, \lambda, \frac{z}{c\tau + d}, \gamma\tau) = (c\tau + d)^k P_k((\mu, \lambda)\gamma, z, \tau).$$

for all $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$.

In order to define Q -functions we need to recall the Bernoulli polynomials $B_r(x) \in \mathbb{Q}[x]$ which are determined by

$$\frac{te^{tx}}{(e^t - 1)} = \sum_{r=0}^{\infty} \frac{B_r(x)t^r}{r!}.$$

For example, $B_0(x) = 1, B_1(x) = x - \frac{1}{2}, B_2(x) = x^2 - x + \frac{1}{6}$.

For $(u, \lambda) = (e^{\frac{2\pi i j}{M}}, e^{\frac{2\pi i l}{N}})$ and $(u, \lambda) \neq (1, 1)$, when $k \geq 1$ and $k \in \mathbb{Z}$, we define

$$\begin{aligned} Q_k(\mu, \lambda, q_\tau) &= Q_k(\mu, \lambda, \tau) \\ &= \frac{1}{(k-1)!} \sum_{n \geq 0} \frac{\lambda(n + j/M)^{k-1} q_\tau^{n+j/M}}{1 - \lambda q_\tau^{n+j/M}} \\ &\quad + \frac{(-1)^k}{(k-1)!} \sum_{n \geq 1} \frac{\lambda^{-1}(n - j/M)^{k-1} q_\tau^{n-j/M}}{1 - \lambda^{-1} q_\tau^{n-j/M}} - \frac{B_k(j/M)}{k!}. \end{aligned} \quad (2.4)$$

Here $(n + j/M)^{k-1} = 1$ if $n = 0, j = 0$ and $k = 1$. Similarly, $(n - j/M)^{k-1} = 1$ if $n = 1, j = M$ and $k = 1$. We also define

$$Q_0(\mu, \lambda, \tau) = -1. \quad (2.5)$$

Here are the transformation properties of Q -functions (see Theorem 4.6 of [DLM3]).

Theorem 2.2 *If $k \geq 0$ then $Q_k(\mu, \lambda, \tau)$ is a holomorphic modular form of weight k . If $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ it satisfies*

$$Q_k(\mu, \lambda, \gamma\tau) = (c\tau + d)^k Q_k((\mu, \lambda)\gamma, \tau).$$

For the further discussion we also define $\bar{P}$ -functions. Again we take $\mu = e^{2\pi ij/M}$ and $\lambda = e^{2\pi il/N}$. Set for $k \geq 1$,

$$\bar{P}_k(\mu, \lambda, z, q_\tau) = \bar{P}_k(\mu, \lambda, z, \tau) = \frac{1}{(k-1)!} \sum'_{n \in j/M + \mathbb{Z}} \frac{n^{k-1} z^n}{1 - \lambda q_\tau^n} \quad (2.6)$$

with

$$\bar{P}_0 = 0. \quad (2.7)$$

Then we have (see Proposition 4.9 of [DLM3]):

Proposition 2.3 *If $m \in \mathbb{Z}$, $\mu = e^{2\pi ij/M}$, $k \geq 0$ and $(\mu, \lambda) \neq (1, 1)$, then*

$$\begin{aligned} & Q_k(\mu, \lambda, \tau) + \frac{1}{k!} B_k(1 - m + j/M) \\ &= \text{Res}_z \left(\sum_{s \geq 0} \frac{z_1^s}{z_1^{s+1}} z_1^{m-j/M} z^{-m+j/M} \bar{P}_k(\mu, \lambda, \frac{z_1}{z}, \tau) \right) \\ &+ \text{Res}_z \left(\lambda \sum_{s \geq 0} \frac{z^s}{z_1^{s+1}} z_1^{m-j/M} z^{-m+j/M} \bar{P}_k(\mu, \lambda, \frac{z_1 q_\tau}{z}, \tau) \right). \end{aligned}$$

We also recall the Eisenstein series of weight k for even $k \geq 3$:

$$\begin{aligned} G_k(\tau) &= \sum'_{m_1, m_2 \in \mathbb{Z}} \frac{1}{(m_1 \tau + m_2)^k} \\ &= (2\pi i)^k \left(\frac{-B_k(0)}{k!} + \frac{2}{(k-1)!} \sum_{n=1}^{\infty} \sigma_{k-1}(n) q^n \right) \end{aligned} \quad (2.8)$$

where $\sigma_{k-1}(n) = \sum_{d|n} d^{k-1}$. For this range of values of k , $G_k(\tau)$ is a modular form on $SL(2, \mathbb{Z})$ of weight k . We will use of the normalized Eisenstein series

$$E_k(\tau) = \frac{1}{(2\pi i)^k} G_k(\tau) = \frac{-B_k(0)}{k!} + \frac{2}{(k-1)!} \sum_{n=1}^{\infty} \sigma_{k-1}(n) q^n \quad (2.9)$$

for even $k \geq 2$. Note that $E_2(\tau)$ is not a modular form but it is a well defined holomorphic function in $0 < |q| < 1$.

3 Vertex operator superalgebras

In this section we recall the definition of vertex operator superalgebra (cf. [B1], [DL], [FLM3]). We also study another vertex operator superalgebra structure on torus following [Z] to give the modularity result in later sections.

Let $V = V_0 \oplus V_1$ be any $\mathbb{Z}_2$ -graded vector space. The element in V_0 (resp. V_1) is called even (resp. odd). For any $v \in V_i$ with $i = 0, 1$ we define $\tilde{v} = i$. For convenience, we let $\epsilon_{u,v} = (-1)^{\tilde{u}\tilde{v}}$ and $\epsilon_v = (-1)^{\tilde{v}}$.

Definition 3.1 A vertex operator superalgebra (VOSA) is a quadruple $(V, Y, \mathbf{1}, \omega)$, where $V = V_0 \oplus V_1$ is a $\mathbb{Z}_2$ -graded vector space, $\mathbf{1} \in V_0$ is the vacuum of V , $\omega \in V_2$ is called conformal vector of V , and Y is a linear map

$$\begin{aligned} V &\rightarrow (\text{End } V)[[z, z^{-1}]], \\ v &\mapsto Y(v, z) = \sum_{n \in \mathbb{Z}} v(n) z^{-n-1} \quad (v(n) \in \text{End } V). \end{aligned}$$

satisfying the following axioms for $u, v \in V$:

- (i) For any $u, v \in V$, $u(n)v = 0$ for sufficiently large n ,
- (ii) $Y(\mathbf{1}, z) = Id_V$,
- (iii) $Y(v, z)\mathbf{1} = v + \sum_{n \geq 2} v(-n)\mathbf{1}z^{n-1}$,
- (iv) The component operators of $Y(\omega, z) = \sum_{n \in \mathbb{Z}} L(n)z^{-n-2}$ satisfy the Virasoro algebra relation with central charge $c \in \mathbb{C}$:

$$[L(m), L(n)] = (m - n)L(m + n) + \frac{1}{12}(m^3 - m)\delta_{m+n,0}c, \quad (3.1)$$

and

$$L(0)|_{V_n} = n, \quad n \in \mathbb{Z} \quad (3.2)$$

$$\frac{d}{dz}Y(v, z) = Y(L(-1)v, z), \quad (3.3)$$

- (v) The Jacobi identity for $\mathbb{Z}_2$ -homogeneous $u, v \in V$ holds,

$$\begin{aligned} z_0^{-1}\delta\left(\frac{z_1 - z_2}{z_0}\right)Y(u, z_1)Y(v, z_2) - \epsilon_{u,v}z_0^{-1}\delta\left(\frac{z_2 - z_1}{-z_0}\right)Y(v, z_2)Y(u, z_1) \\ = z_2^{-1}\delta\left(\frac{z_1 - z_0}{z_2}\right)Y(Y(u, z_0)v, z_2) \end{aligned} \quad (3.4)$$

where $\delta(z) = \sum_{n \in \mathbb{Z}} z^n$ and $(z_i - z_j)^n$ is expanded as a formal power series in z_j . Throughout the paper, z_0, z_1, z_2 , ect. are independent commuting formal variables.

As in [Z], we also define a vertex operator superalgebra on torus associated to V . Set $\tilde{\omega} = \omega - \frac{c}{24}$ and

$$Y[v, z] = Y(v, e^z - 1)e^{z\omega v} = \sum_{n \in \mathbb{Z}} v[n]z^{-n-1} \quad (3.5)$$

for homogeneous $v \in V$. Following the proof of Theorem 4.21 of [Z] we have

Theorem 3.2 $(V, Y[\cdot], \mathbf{1}, \tilde{\omega})$ is a vertex operator superalgebra.

Here we compute $v[m]$. Using a change of variable we calculate that

$$\begin{aligned} v[m] &= \text{Res}_z Y[v, z] z^m \\ &= \text{Res}_z Y[v, \log(1+z)] (\log(1+z))^m (1+z)^{-1} \end{aligned}$$

i.e.,

$$v[m] = \text{Res}_z Y(v, z) (\log(1+z))^m (1+z)^{\text{wt}v-1}. \quad (3.6)$$

For integers i, m, p with $i, m \geq 0$, define numbers $c(p, i, m)$ by two equivalent ways:

$$\binom{p-1+z}{i} = \sum_{m=0}^{\infty} c(p, i, m) z^m. \quad (3.7)$$

It is easy to see that

$$m! \sum_{i=m}^{\infty} c(p, i, m) z^i = (\log(1+z))^m (1+z)^{p-1} \quad (3.8)$$

where,

$$\log(1+z) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} z^n. \quad (3.9)$$

If $m \geq 0$ then

$$v[m] = m! \sum_{i=m}^{\infty} c(\text{wt}v, i, m) v(i). \quad (3.10)$$

In particular,

$$v[0] = \sum_{i=0}^{\infty} \binom{\text{wt}v-1}{i} v(i). \quad (3.11)$$

Write

$$Y[\tilde{\omega}, z] = \sum_{n \in \mathbb{Z}} L[n] z^{-n-2}. \quad (3.12)$$

One can verify that

$$L[-2] = \omega[-1] - \frac{c}{24} \quad (3.13)$$

$$L[-1] = L(-1) + L(0) \quad (3.14)$$

$$L[0] = L(0) + \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n(n+1)} L(n). \quad (3.15)$$

If $v \in V$ is homogeneous in $(V, Y[\cdot], \mathbf{1}, \tilde{\omega})$, we denote its conformal weight by $\text{wt}[v]$.

Definition 3.3 An automorphism g of vertex operator algebra V is a linear automorphism of V preserving $\mathbf{1}$ and ω such that the actions of g and $Y(v, z)$ on V are compatible in the sense that

$$gY(v, z)g^{-1} = Y(gv, z)$$

for $v \in V$.

Note that an automorphism preserves each homogeneous space V_n and thus each V_i . There is a special automorphism σ of V with $\sigma|V_i = (-1)^i$ associated to the superspace structure of V . Then σ is a central element in the full automorphism group $\text{Aut}(V)$. As we will see later that σ plays a fundamental role in formulating the modular invariance of trace functions.

4 Twisted Modules for VOSA

The twisted modules are the main objects in this section. In the case of vertex operator algebras, the twisted modules are studied in [D], [DLM2], [FFR], [FLM2], [FLM3], [L]. We will follow [DLM2] and [DZ] in this section. In particular, we will introduce the notion of weak, admissible and ordinary twisted modules. We also present the associative algebra $A_g(V)$ associated to a vertex operator algebra V and a finite order automorphism g and related consequences.

Let $(V, \mathbf{1}, \omega, Y)$ be a VOSA and g an automorphism of V of finite order T . Let T' be the order of $g\sigma$. Then we have the following eigenspace decompositions:

$$V = \bigoplus_{r \in \mathbb{Z}/T'\mathbb{Z}} V^{r*} \quad (4.1)$$

$$V = \bigoplus_{r \in \mathbb{Z}/T\mathbb{Z}} V^r \quad (4.2)$$

where $V^{r*} = \{v \in V | g\sigma v = e^{-2\pi i r/T'} v\}$ and $V^r = \{v \in V | gv = e^{-2\pi i r/T} v\}$. We now can define various notions of g -twisted modules (cf. [DZ]).

Definition 4.1 *A weak g -twisted V -module is a $\mathbb{C}$ -linear space M equipped with a linear map $V \rightarrow (\text{End} M)[[z^{1/T}, z^{-1/T}]]$, $v \mapsto Y_M(v, z) = \sum_{n \in \mathbb{Q}} v(n) z^{-n-1}$ satisfying:*

- (1) *For $v \in V, w \in M$, $v(m)w = 0$ if $m \gg 0$,*
- (2) *$Y_M(\mathbf{1}, z) = Id_V$,*
- (3) *For $v \in V^r$, and $0 \leq r \leq T-1$,*

$$Y_M(v, z) = \sum_{n \in r/T + \mathbb{Z}} v(n) z^{-n-1} \quad \text{for } v \in V^r,$$

- (4) *The twisted Jacobi identity holds:*

$$\begin{aligned} z_0^{-1} \delta\left(\frac{z_1 - z_2}{z_0}\right) Y_M(u, z_1) Y_M(v, z_2) - \epsilon_{u,v} z_0^{-1} \delta\left(\frac{z_2 - z_1}{-z_0}\right) Y_M(v, z_2) Y_M(u, z_1) \\ = z_2^{-1} \left(\frac{z_1 - z_0}{z_2}\right)^{-r/T} \delta\left(\frac{z_1 - z_0}{z_2}\right) Y_M(Y(u, z_0)v, z_2) \end{aligned} \quad (4.3)$$

if $u \in V^r$.

Note that (4.3) is equivalent to the following:

$$\begin{aligned} & \text{Res}_{z_1} Y_M(u, z_1) Y_M(v, z_2) z_1^{r/T} (z_1 - z_2)^m z_2^n \\ & - \epsilon_{u,v} \text{Res}_{z_1} Y_M(v, z_2) Y_M(u, z_1) z_1^{r/T} (z_1 - z_2)^m z_2^n \\ & = \text{Res}_{z_1 - z_2} Y_M(Y(u, z_1 - z_2)v, z_2) \iota_{z_2, z_1 - z_2} z_1^{r/T} (z_1 - z_2)^m z_2^n \end{aligned} \quad (4.4)$$

for $m \in \mathbb{Z}, n \in \mathbb{C}$, where $\iota_{z_2, z_1 - z_2} z_1^{r/T} = \sum_{m \geq 0} \binom{r/T}{m} z_2^{r/T - m} (z_1 - z_2)^m$.

Set

$$Y_M(\omega, z) = \sum_{n \in \mathbb{Z}} L(n) z^{-n-2}.$$

Then $Y_M(L(-1)z, v) = \frac{d}{dz} Y_M(v, z)$ for $v \in V$ and the operators $L(n)$ satisfy the Virasoro algebra relation with central charge c (cf. [DLM1], [DZ]).

Definition 4.2 *An admissible g -twisted V -module is a weak g -twisted V -module M which carries a $\frac{1}{T'}$ $\mathbb{Z}$ -grading*

$$M = \bigoplus_{0 \leq n \in \frac{1}{T'}\mathbb{Z}} M(n) \quad (4.5)$$

such that

$$v(m)M(n) \subseteq M(n + \text{wt}v - m - 1) \quad (4.6)$$

for homogeneous $v \in V$. We may and do assume that $M(0) \neq 0$ if $M \neq 0$.

Note that an admissible V -module M is $\frac{1}{T'}\mathbb{Z}$ -graded instead of $\frac{1}{T}\mathbb{Z}$ -graded as V is $\frac{1}{2}\mathbb{Z}$ -graded.

Definition 4.3 *An (ordinary) g -twisted V -module M is a $\mathbb{C}$ -graded weak g -twisted V -module*

$$M = \coprod_{\lambda \in \mathbb{C}} M_\lambda \quad (4.7)$$

where $M_\lambda = \{w \in M \mid L(0)w = \lambda w\}$ such that $\dim M_\lambda$ is finite and for fixed λ , $M_{\frac{n}{T'} + \lambda} = 0$ for all small enough integers n .

It $g = 1$ we have the notion of weak, admissible ordinary V -modules. It is not hard to prove that any ordinary g -twisted V -module is admissible.

If M is a simple g -twisted V -module, then

$$M = \bigoplus_{n=0}^{\infty} M_{\lambda + n/T'} \quad (4.8)$$

for some $\lambda \in \mathbb{C}$ such that $M_\lambda \neq 0$ (cf. [Z]). We define λ as the *conformal weight* of M .

Definition 4.4 (1) *A VOSA V is called rational for an automorphism g of finite order if the category of admissible modules is semisimple.*

(2) *V is called g -regular if any weak g -twisted V -module is a direct sum of irreducible ordinary g -twisted V -modules.*

It is clear that a g -regular vertex operator superalgebra is g -rational.

The following theorem is proved in [DZ].

Theorem 4.5 *Let V be a VOSA and g an automorphism of V of finite order. Assume that V is g -rational. Then*

- (1) *There are only finitely many irreducible admissible g -twisted V -modules up to isomorphism.*
- (2) *Each irreducible admissible g -twisted V -module is ordinary.*

Now we review the $A_g(V)$ theory [DZ]. Let V^{r*} be as in (4.1). For $u \in V^{r*}$ and $v \in V$ we define

$$u \circ_g v = \text{Res}_z \frac{(1+z)^{\text{wt}u-1+\delta_r+\frac{r}{T}}}{z^{1+\delta_r}} Y(u, z)v \quad (4.9)$$

$$u *_g v = \begin{cases} \text{Res}_z(Y(u, z) \frac{(1+z)^{\text{wt}u}}{z} v) & \text{if } r = 0 \\ 0 & \text{if } r > 0. \end{cases} \quad (4.10)$$

where $\delta_r = 1$ if $r = 0$ and $\delta_r = 0$ if $r \neq 0$. Extend $\circ_g$ and $*_g$ to bilinear products on V . We let $O_g(V)$ be the linear span of all $u \circ_g v$. Then the quotient $A_g(V) = V/O_g(V)$ is an associative algebra with respect to $*_g$.

Let M be a weak σg -twisted V -module. Define

$$\Omega(M) = \{w \in M | u(\text{wt}u - 1 + n)w = 0, u \in V, n > 0\}.$$

For homogeneous $u \in V$ we set

$$o(u) = u(\text{wt}u - 1) \quad (4.11)$$

We have the following Theorem (see [DZ]):

Theorem 4.6 *Let V be a VOSA together with an automorphism g of finite order, and M a weak g -twisted V -module. Then*

- (1) *$\Omega(M)$ is an $A_g(V)$ -module such that $v + O_g(V)$ acts as $o(v)$.*
- (2) *If $M = \sum_{n \geq 0} M(n/T)$ is an admissible g -twisted V -module with $M(0) \neq 0$, then $M(0) \subset \Omega(M)$ is an $A_g(V)$ -submodule. Moreover, M is irreducible if and only if $M(0) = \Omega(M)$ and $M(0)$ is a simple $A_g(V)$ -module.*
- (3) *The map $M \rightarrow M(0)$ gives a 1-1 correspondence between the irreducible admissible g -twisted V -modules and simple $A_g(V)$ -modules.*
- (4) *If V is g -rational then $A_g(V)$ is a finite dimensional semisimple associative algebra.*

Another important concept is the C_2 -cofinite condition (cf. [Z]). Set

$$C_2(V) = \{u(-2)v | u, v \in V\}. \quad (4.12)$$

V is called C_2 -cofinite if $C_2(V)$ has finite codimension.

It is proved in [Li] and [ABD] that a vertex operator algebra V is regular if and only if V is C_2 -cofinite and rational. Using the the same proof we have

Theorem 4.7 *Assume that V is C_2 -cofinite. Then V is g -rational if and only if V is g -regular.*

Here is another consequence of the C_2 -cofiniteness.

Proposition 4.8 *Suppose that V satisfies condition C_2 , and let g be an automorphism of V of finite order. Then*

- (1) *The algebra $A_g(V)$ has finite dimension.*
- (2) *If $A_g(V) \neq 0$ then V has a simple g -twisted V -module.*

The proof of (1) is similar to that of Proposition 3.6 in [DLM3] and (2) follows from Theorem 4.6 (3).

5 1-point functions on the torus

We first introduce some notations which will be used in this section. V again is a VOSA and $g, h \in \text{Aut}(V)$ such that $gh = hg$. Let $o(g) = T, o(h) = T_1$ be finite.

Let A be the subgroup of $\text{Aut}(V)$ generated by $g\sigma$ and $h\sigma$, and $N = \text{lcm}(T, T_1)$ be the exponent of A . Let $\Gamma(T, T_1)$ be the subgroup of matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in $SL(2, \mathbb{Z})$ satisfying $a \equiv d \equiv 1 \pmod{N}$, $b \equiv 0 \pmod{T}$, $c \equiv 0 \pmod{T_1}$, and $M(T, T_1)$ be the ring of holomorphic modular forms on $\Gamma(T, T_1)$ with natural gradation $M(T, T_1) = \bigoplus_{k \geq 0} M_k(T, T_1)$, where $M_k(T, T_1)$ is the space of forms of weight k . From the Lemma 5.1 in [DLM3], $M(T, T_1)$ is a Noetherian ring which contains each $E_{2k}(\tau)$, $k \geq 2$, and each $Q_k(\mu, \lambda, \tau)$, $k \geq 0$ for μ, λ a T -th., resp. T_1 -th. root of unity.

Let $V(T, T_1) = M(T, T_1) \otimes_{\mathbb{C}} V$. For $v \in V$ with $gv = \mu^{-1}v, hv = \lambda^{-1}v$, we can define a $M(T, T_1)$ -submodule of $V(T, T_1)$, say $O(g, h)$ which consists of the following elements:

$$v[0]w, w \in V, (\mu, \lambda) = (1, 1) \tag{5.1}$$

$$v[-2]w + \sum_{k=2}^{\infty} (2k-1)E_{2k}(\tau) \otimes v[2k-2]w, (\mu, \lambda) = (1, 1) \tag{5.2}$$

$$v, (\tilde{v}, \mu, \lambda) \neq (1, 1, 1) \tag{5.3}$$

$$\sum_{k=0}^{\infty} Q_k(\mu, \lambda, \tau) \otimes v[k-1]w, (\mu, \lambda) \neq (1, 1). \tag{5.4}$$

We have the following Lemmas as in [DLM3].

Lemma 5.1 *Suppose V satisfies C_2 -cofinite condition. Then $V(T, T_1)/O(g, h)$ is a finitely-generated $M(T, T_1)$ -module.*

Lemma 5.2 *If V is C_2 -cofinite, then for $v \in V$, there is $m \in \mathbb{N}$ and elements $r_i(\tau) \in M(T, T_1)$, $0 \leq i \leq m-1$, such that*

$$L[-2]^m v + \sum_{i=0}^{m-1} r_i(\tau) \otimes L[-2]^i v \in O(g, h). \tag{5.5}$$

Definition 5.3 *The space of (g, h) 1-point functions $\mathcal{C}(g, h)$ is a $\mathbb{C}$ -linear space consisting of functions*

$$S : V(T, T_1) \times \mathfrak{h} \rightarrow \mathbb{C}$$

satisfying the following conditions:

- (1) $S(v, \tau)$ is holomorphic in τ for $v \in V(T, T_1)$.
- (2) $S(v, \tau)$ is $\mathbb{C}$ linear in v and for $f \in M(T, T_1)$, $v \in V$,

$$S(f \otimes v, \tau) = f(\tau)S(v, \tau)$$

- (3) $S(v, \tau) = 0$ if $v \in O(g, h)$.
- (4) If $v \in V$ with $\sigma v = gv = hv = v$, then

$$S(L[-2]v, \tau) = \partial S(v, \tau) + \sum_{l=2}^{\infty} E_{2l}(\tau) S(L[2l-2]v, \tau). \quad (5.6)$$

Here ∂S is the operator which is linear in v and satisfies

$$\partial S(v, \tau) = \partial_k S(v, \tau) = \frac{1}{2\pi i} \frac{d}{d\tau} S(v, \tau) + kE_2(\tau)S(v, \tau) \quad (5.7)$$

for $v \in V_{[k]}$.

The following result – modular transformation is the main property of 1-point functions.

Theorem 5.4 *For $S \in \mathcal{C}(g, h)$ and $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$, we define*

$$S|_{\gamma}(v, \tau) = S|_k \gamma(v, \tau) = (c\tau + d)^{-k} S(v, \gamma\tau) \quad (5.8)$$

for $v \in V_{[k]}$, and extend linearly. Then $S|_{\gamma} \in \mathcal{C}((g, h)\gamma)$.

This theorem is the ‘super’ analogue of Theorem 5.4 in [DLM3] and the proof is the same.

Now we study the q -expansion of $S(v, \tau)$ for $S \in \mathcal{C}(g, h)$ and $v \in V$. As argued in [DLM3], the Frobenius-Fuchs theory tells us that $S(v, \tau)$ may be expressed in the following form: for some $p \geq 0$,

$$S(v, \tau) = \sum_{i=0}^p (\log q_{\frac{1}{T}})^i S_i(v, \tau) \quad (5.9)$$

where

$$S_i(v, \tau) = \sum_{j=1}^{b(i)} q^{\lambda_{i,j}} S_{i,j}(v, \tau) \quad (5.10)$$

$$S_{i,j}(v, \tau) = \sum_{n=0}^{\infty} a_{i,j,n}(v) q^{n/T} \quad (5.11)$$

are holomorphic on the upper half-plane, and

$$\lambda_{i,j_1} \not\equiv \lambda_{i,j_2} \pmod{\frac{1}{T}\mathbb{Z}} \quad (5.12)$$

for $j_1 \neq j_2$ where $q_{\frac{1}{T}} = e^{2\pi i\tau/T}$ and $\lambda_{i,j} \in \mathbb{C}$. Moreover, p is bounded independently of v .

It is important to point out that the 1-point functions on the torus can be defined formally regarding q as a formal variable. That is, we identify elements of $M(T, T_1)$ with their Fourier expansions at ∞ , which lie in the ring of formal power series $\mathbb{C}[[q_{\frac{1}{T}}]]$. Similarly, the functions $E_{2k}(\tau)$, $k \geq 1$, are considered to lie in $\mathbb{C}[[q]]$. The operator $\frac{1}{2\pi i} \frac{d}{d\tau}$ acts on $\mathbb{C}[[q_{\frac{1}{T}}]]$ via the identification $\frac{1}{2\pi i} \frac{d}{d\tau} = q \frac{d}{dq}$.

Then a *formal* (g, h) 1-point function is a map

$$S : V(T, T_1) \rightarrow P$$

where P is the space of formal power series of the form

$$q^\lambda \sum_{n=0}^{\infty} a_n q^{n/T} \quad (5.13)$$

for some $\lambda \in \mathbb{C}$, and it satisfies the formal analogues of (2)-(4) in the definition of (g, h) 1-point functions. Then as in [DLM3], each $S(v, q)$ converges to a holomorphic function. we have

Theorem 5.5 *Assume that V is C_2 -cofinite. Then any formal (g, h) 1-point function S defines an element of $\mathcal{C}(g, h)$, also denoted by S , via the identification*

$$S(v, \tau) = S(v, q), q = q_\tau = e^{2\pi i\tau}. \quad (5.14)$$

6 Trace functions

The goal in this section is to construct (g, h) 1-point functions which are essentially the graded $h\sigma$ -trace function on $g\sigma$ -twisted V -modules.

Let $M = (M, Y_M)$ be a $g\sigma$ -twisted V -module and $k \in \text{Aut}(V)$. We can define a $k(\sigma g)k^{-1}$ -twisted V -module $(k \circ M, Y_{k \circ M})$ so that $k \circ M = M$ as vector spaces and

$$Y_{k \circ M}(v, z) = Y_M(k^{-1}v, z).$$

M is called k -stable if $k \circ M$ and M are isomorphic.

Recall from Section 4 that if M is a simple $g\sigma$ -module then there exists a complex number λ such that

$$M = \bigoplus_{n=0}^{\infty} M_{\lambda + \frac{n}{T}} \quad (6.1)$$

(see (4.8)) where T is the order of g . In particular, if $g = \sigma$ then

$$M = \bigoplus_{n=0}^{\infty} M_{\lambda + \frac{n}{2}}. \quad (6.2)$$

Let $h \in \text{Aut}(V)$ as before and assume that $h\sigma \circ M$ and M are isomorphic. Then g, h commute and there is a linear map $\phi(h\sigma) : M \rightarrow M$ such that

$$\phi(\sigma h)Y_M(v, z)\phi(\sigma h)^{-1} = Y_M((\sigma h)v, z) \quad (6.3)$$

for all $v \in V$.

Lemma 6.1 *If M is a simple $g\sigma$ -twisted V -module then $g \circ M$ and M are isomorphic. In particular, any simple V -module M and $\sigma \circ M$ are isomorphic.*

Proof: We define a map $\phi(g)$ on M such that $\phi(g)|_{M_{\lambda+n/T}} = e^{2\pi i n/T}$. Let $o(g\sigma) = T'$ and $v \in V^{r*}$, $v \in V^s$ (see (4.1)-(4.2)). It is easy to verify that $\text{wt}v - \frac{r}{T'}$ and $\frac{s}{T}$ are congruent modulo $\mathbb{Z}$. We immediately have $\phi(g)Y_M(v, z)\phi(g)^{-1} = Y_M(gv, z)$ for all $v \in V$. That is, M and $g \circ M$ are isomorphic. $\square$

We remark that this result is different from a similar result in the case of vertex operator algebra where a simple g -twisted module M and $g \circ M$ are always isomorphic. But this is not true in the super case.

Now we introduce the function T which is linear in $v \in V$, and define for homogeneous $v \in V$ as follows:

$$T(v) = T_M(v, (g, h), q) = z^{\text{wt}v} \text{tr}_M Y_M(v, z) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \quad (6.4)$$

Here c is the central charge of V . Note that for $m \in \frac{1}{T}$, $v(m)$ maps M_μ to $M_{\mu+\text{wt}v-m-1}$. So we can write $T(v)$ as follows:

$$T(v) = q^{\lambda - \frac{c}{24}} \sum_{n=0}^{\infty} \text{tr}_{M_{\lambda+\frac{n}{T}}} o(v) \phi(\sigma h) q^{\frac{n}{T}} = \text{tr}_M o(v) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \quad (6.5)$$

Here is the main result in this section.

Theorem 6.2 *Let V be C_2 -cofinite and $g, h \in \text{Aut}(V)$ have finite orders. Let M be a simple $g\sigma$ -twisted V -module such that $h \circ M, \sigma \circ M$ and M are isomorphic. Then $T \in \mathcal{C}(g, h)$.*

By Theorem 5.5 it is enough to prove that $T(v)$ defines a formal (g, h) 1-point function. Clearly, $T(v)$ has the shape of (5.13). So we need to prove that $T(v)$ satisfies (2)-(4) in definition 5.3. But (2) is clear, we are going to prove (3) and (4).

Fix $v \in V$ such that $gv = u^{-1}v$, $hv = \lambda^{-1}v$ and $v_1 \in V$.

Lemma 6.3 *$T(v) = 0$ for $(\epsilon_v, \mu, \lambda) \neq (1, 1, 1)$.*

Proof: Let $k \in \{g, h, \sigma\}$. Since $k \circ M$ and M are isomorphic (see Lemma 6.1), then there is a linear isomorphism $\phi(k) : M \rightarrow M$ such that $\phi(k)Y_M(v, z)\phi(k)^{-1} = Y_M(kv, z)$ for all $v \in V$. Normalizing $\phi(k)$ we can assume that $\phi(k)^t = 1$ where t is the order of k . So we can decompose M into a direct sum of eigenspaces. Since g, h, σ commute with each other, $\phi(h\sigma)$ preserves each eigenspace. Since v is an eigenvector for k with eigenvalue different from 1, $o(v)$ moves one eigenspace of $\phi(k)$ to a different eigenspace. As a result, the trace of $o(v)\phi(h\sigma)$ on any homogeneous subspace $M_{\lambda+n/T}$ is zero. The proof is complete. $\square$

Lemma 6.4 $T(v[0]v_1) = 0$ for $(u, \lambda) = (1, 1)$.

Proof: Consider

$$\begin{aligned}
& \text{tr}_M[v(\text{wt}v - 1), Y_M(v_1, z)]\phi(\sigma h)q^{L(0)-c/24} \\
&= \text{tr}_M v(\text{wt}v - 1)Y_M(v_1, z)\phi(\sigma h)q^{L(0)-c/24} - \epsilon_{v,v_1} \text{tr}_M Y_M(v_1, z)v(\text{wt}v - 1)\phi(\sigma h)q^{L(0)-c/24} \\
&= \text{tr}_M v(\text{wt}v - 1)Y_M(v_1, z)\phi(\sigma h)q^{L(0)-c/24} - \epsilon_{v,v_1} \epsilon_v \text{tr}_M Y_M(v_1, z)\phi(\sigma h)q^{L(0)-c/24} v(\text{wt}v - 1) \\
&= (1 - \epsilon_{v,v_1} \epsilon_v) \text{tr}_M v(\text{wt}v - 1)Y_M(v_1, z)\phi(\sigma h)q^{L(0)-c/24}.
\end{aligned}$$

On the other hand

$$\begin{aligned}
& \text{tr}_M[v(\text{wt}v - 1), Y_M(v_1, z)]\phi(\sigma h)q^{L(0)-c/24} \\
&= \text{tr}_M \sum_{i=0}^{\infty} \binom{\text{wt}v - 1}{i} z^{\text{wt}v - 1 - i} Y_M(v(i)v_1, z)\phi(\sigma h)q^{L(0)-c/24} \\
&= z^{-\text{wt}v} T(v[0]v_1)
\end{aligned}$$

Thus if $v \in V_{\bar{0}}$, or both v and v_1 are in $V_{\bar{1}}$, then $\epsilon_{v,v_1} \epsilon_v = 1$ and $T(v[0]v_1) = 0$. If $v \in V_{\bar{1}}$, and $v_1 \in V_{\bar{0}}$, then $v[0]v_1 \in V_{\bar{1}}$. By Lemma 6.3, we also have $T(v[0]v_1) = 0$. $\square$

Lemma 6.5

$$\sum_{k=0}^{\infty} Q_k(\mu, \lambda, \tau) \otimes T(v[k-1]v_1) = 0$$

for $(\mu, \lambda) \neq (1, 1)$.

In order to prove Lemma 6.5, we first define 2-point correlation functions. These are multi-linear functions $T(v_1, v_2)$ defined for $v_1, v_2 \in V$ homogeneous via

$$\begin{aligned}
T(v_1, v_2) &= T((v_1, z_1), (v_2, z_2), (g, h), q) \\
&= z_1^{\text{wt}v_1} z_2^{\text{wt}v_2} \text{tr} Y_M(v_1, z_1) Y_M(v_2, z_2) \phi(\sigma h) q^{L(0)-c/24}.
\end{aligned} \tag{6.6}$$

We need the following sublemma.

Sublemma 6.6 *Let $v, v_1 \in V$ be homogeneous with $gv = \mu^{-1}v$, $hv = \lambda^{-1}v$ and $(\mu, \lambda) \neq (1, 1)$. Then*

$$T(v, v_1) = \sum_{k=1}^{\infty} \bar{P}_k(\mu, \lambda, \frac{z_1}{z}, q) T(v[k-1]v_1) \tag{6.7}$$

$$T(v_1, v) = (-1)^{\tilde{v}} \lambda \sum_{k=1}^{\infty} \bar{P}_k(\mu, \lambda, \frac{z_1}{z} q, q) T(v[k-1]v_1) \tag{6.8}$$

where $\bar{P}_k$ is as in (2.6).

Proof: First we have the following identities:

$$\begin{aligned} & (1 - \epsilon_{v,v_1} \epsilon_v \lambda q^k) \text{tr}_M v(\text{wt}v - 1 + k) Y_M(v_1, z_1) \phi(\sigma h) q^{L(0)} \\ &= \sum_{i=0}^{\infty} \binom{\text{wt}v - 1 + k}{i} z_1^{\text{wt}v - 1 + k - i} \text{tr}_M Y_M(v(i)v_1, z_1) \phi(\sigma h) q^{L(0)} \end{aligned} \quad (6.9)$$

$$\begin{aligned} & (1 - \epsilon_{v,v_1} \epsilon_v \lambda q^k) \text{tr}_M Y_M(v_1, z_1) v(\text{wt}v - 1 + k) \phi(\sigma h) q^{L(0)} \\ &= \epsilon_v \lambda q^k \sum_{i=0}^{\infty} \binom{\text{wt}v - 1 + k}{i} z_1^{\text{wt}v - 1 + k - i} \text{tr}_M Y_M(v(i)v_1, z_1) \phi(\sigma h) q^{L(0)}. \end{aligned} \quad (6.10)$$

The proofs of these identities are similar to that given in Lemma 8.4 of [DLM3] by noting the following

$$\begin{aligned} \text{tr}_M v(\text{wt}v - 1 + k) Y_M(v_1, z_1) \phi(\sigma h) q^{L(0)} &= \text{tr}_M [v(\text{wt}v - 1 + k), Y_M(v_1, z_1)] \phi(\sigma h) q^{L(0)} \\ &+ \epsilon_{v,v_1} \text{tr}_M Y_M(v_1, z_1) v(\text{wt}v - 1 + k) \phi(\sigma h) q^{L(0)}, \end{aligned} \quad (6.11)$$

$$v(\text{wt}v - 1 + k) \phi(\sigma h) = \epsilon_v \lambda \phi(\sigma h) v(\text{wt}v - 1 + k),$$

and

$$\text{tr}_M Y_M(v_1, z_1) v(\text{wt}v - 1 + k) \phi(\sigma h) q^{L(0)} = \epsilon_v \lambda q^k \text{tr}_M v(\text{wt}v - 1 + k) Y_M(v_1, z_1) \phi(\sigma h) q^{L(0)}. \quad (6.12)$$

Use identities (6.9), (6.10) and the same proof of Theorem 8.4 of [DLM3] to obtain:

$$\begin{aligned} T(v, v_1) &= \sum_{k=1}^{\infty} \bar{P}_k(\mu, \epsilon_{v,v_1} \epsilon_v \lambda, \frac{z_1}{z}, q) T(v[k-1]v_1) \\ T(v_1, v) &= \epsilon_v \lambda^{-1} \sum_{k=1}^{\infty} \bar{P}_k(\mu, \epsilon_{v,v_1} \epsilon_v \lambda, \frac{z_1}{z} q, q) T(v[k-1]v_1). \end{aligned}$$

Again if $\epsilon_{v,v_1} \epsilon_v \neq 1$ then $T(v[k-1]v_1) = 0$ by Lemma 6.3. So in any case we have

$$\begin{aligned} T(v, v_1) &= \sum_{k=1}^{\infty} \bar{P}_k(\mu, \lambda, \frac{z_1}{z}, q) T(v[k-1]v_1) \\ T(v_1, v) &= \epsilon_v \lambda^{-1} \sum_{k=1}^{\infty} \bar{P}_k(\mu, \lambda, \frac{z_1}{z} q, q) T(v[k-1]v_1), \end{aligned}$$

as required. $\square$

We are now in a position to prove Lemma 6.5. Recall the following equality about Bernoulli polynomials from [DLM3], Lemma 8.6.

$$\sum_{k=0}^{\infty} \frac{1}{k!} B_k(1 - \text{wt}v + \frac{r}{T} + \frac{1}{2} \delta_{\bar{v},1}) v[k-1] = \sum_{i=0}^{\infty} \binom{\frac{r}{T} + \frac{1}{2} \delta_{\bar{v},1}}{i} v(i-1). \quad (6.13)$$

Let $m = \text{wt}v - \frac{1}{2}\delta_{\bar{v},1} \in \mathbb{Z}$ in Proposition 2.3. Using (6.13) we have

$$\begin{aligned}
& \sum_{k=0}^{\infty} Q_k(\mu, \lambda, q) T(v[k-1]v_1) \\
&= \sum_{k=1}^{\infty} \text{Res}_z \left((z - z_1)^{-1} z_1^{m - \frac{r}{T} - \frac{1}{2}\delta_{\bar{v},1}} z^{-m + \frac{r}{T} + \frac{1}{2}\delta_{\bar{v},1}} \bar{P}_k(\mu, \lambda, \frac{z_1}{z}, q) \right) T(v[k-1]v_1) \\
&\quad - \lambda \sum_{k=1}^{\infty} \text{Res}_z \left((-z_1 + z)^{-1} z_1^{m - \frac{r}{T} - \frac{1}{2}\delta_{\bar{v},1}} z^{-m + \frac{r}{T} + \frac{1}{2}\delta_{\bar{v},1}} \bar{P}_k(\mu, \lambda, \frac{z_1 q}{z}, q) \right) T(v[k-1]v_1) \\
&\quad - \sum_{i=0}^{\infty} \binom{\frac{r}{T} + \frac{1}{2}\delta_{\bar{v},1}}{i} T(v(i-1)v_1).
\end{aligned}$$

On the other hand, by (4.4),

$$\begin{aligned}
& \sum_{i=0}^{\infty} \binom{\frac{r}{T} + \frac{1}{2}\delta_{\bar{v},1}}{i} T(v(i-1)v_1) \\
&= \sum_{i=0}^{\infty} \binom{\frac{r}{T} + \frac{1}{2}\delta_{\bar{v},1}}{i} z_1^{\text{wt}v + \text{wt}v_1 - i} \text{tr} Y_M(v(i-1)v_1, z_1) \phi(\sigma h) q^{L(0)-c/24} \\
&= \sum_{i=0}^{\infty} \binom{\frac{r}{T} + \frac{1}{2}\delta_{\bar{v},1}}{i} z_1^{\text{wt}v + \text{wt}v_1 - i} \text{Res}_{z=z_1} (z - z_1)^{i-1} \text{tr} Y_M(Y(v, z - z_1)v_1, z_1) \phi(\sigma h) q^{L(0)-c/24} \\
&= \text{Res}_{z=z_1} \iota_{z_1, z-z_1} \left(\frac{z}{z_1} \right)^{\frac{r}{T} + \frac{1}{2}\delta_{\bar{v},1}} (z - z_1)^{-1} z_1^{\text{wt}v + \text{wt}v_1} \text{tr} Y_M(Y(v, z - z_1)v_1, z_1) \phi(\sigma h) q^{L(0)-c/24} \\
&= \text{Res}_z \iota_{z, z_1} (z - z_1)^{-1} \left(\frac{z_1}{z} \right)^{\text{wt}v - \frac{r}{T} - \frac{1}{2}\delta_{\bar{v},1}} T(v, v_1) \\
&\quad - \epsilon_{v, v_1} \text{Res}_z \iota_{z_1, z} (z - z_1)^{-1} \left(\frac{z_1}{z} \right)^{\text{wt}v - \frac{r}{T} - \frac{1}{2}\delta_{\bar{v},1}} T(v_1, v)
\end{aligned}$$

which by Sublemma 6.6 is equal to

$$\begin{aligned}
& \sum_{k=1}^{\infty} \text{Res}_z \iota_{z, z_1} (z - z_1)^{-1} \left(\frac{z_1}{z} \right)^{\text{wt}v - \frac{r}{T} - \frac{1}{2}\delta_{\bar{v},1}} \bar{P}_k(\mu, \lambda, \frac{z_1}{z}, q) T(v[k-1]v_1) \\
&\quad - \epsilon_{v, v_1} \epsilon_v \lambda \sum_{k=1}^{\infty} \text{Res}_z \iota_{z_1, z} (z - z_1)^{-1} \left(\frac{z_1}{z} \right)^{\text{wt}v - \frac{r}{T} - \frac{1}{2}\delta_{\bar{v},1}} \bar{P}_k(\mu, \lambda, \frac{z_1}{z} q, q) T(v[k-1]v_1).
\end{aligned}$$

Again, if $v \in V_{\bar{0}}$ or both $v, v_1 \in V_{\bar{1}}$ then $\epsilon_{v, v_1} \epsilon_v = 1$ and the result follows. If $v \in V_{\bar{1}}$ and $v_1 \in V_{\bar{0}}$, the result follows from Lemma 6.3. $\square$

Lemma 6.7 *If $(\mu, \lambda) = (1, 1)$, then*

$$T(v[-2]v_1) + \sum_{k=2}^{\infty} (2k-1) E_{2k}(\tau) T(v[2k-2]v_1) = 0.$$

Sublemma 6.8 *Let $v, v_1 \in V$ be the same as in Lemma 6.7. Then*

$$T(v, v_1) = \text{tr}_M o(v) o(v_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} + \sum_{k=1}^{\infty} \bar{P}_k(1, 1, \frac{z_1}{z}, q) T(v[k-1]v_1) \quad (6.14)$$

$$T(v_1, v) = \text{tr}_M o(v_1) o(v) \phi(\sigma h) q^{L(0) - \frac{c}{24}} + \epsilon_v \lambda \sum_{k=1}^{\infty} \bar{P}_k(1, 1, \frac{z_1}{z} q, q) T(v[k-1]v_1) \quad (6.15)$$

where $\bar{P}_k$ is defined in (2.6).

Proof: We only prove (6.14) and (6.15) can be proved similarly. By (6.9), (3.7) and (3.10) we have

$$\begin{aligned} T(v, v_1) &= z^{\text{wt}v} z_1^{\text{wt}v_1} \text{tr} Y_M(v, z) Y_M(v_1, z_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \\ &= z^{\text{wt}v} z_1^{\text{wt}v_1} \text{tr} \sum_{k \in \mathbb{Z}} z^{-k - \text{wt}v} v(\text{wt}v - 1 + k) Y_M(v_1, z_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \\ &= z_1^{\text{wt}v_1} \text{tr} o(v) Y_M(v_1, z_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \\ &\quad + z_1^{\text{wt}v_1} \sum_{k \in \mathbb{Z} - \{0\}} \frac{z^{-k}}{1 - \epsilon_{v, v_1} \epsilon_v q^k} \sum_{i=0}^{\infty} \binom{\text{wt}v - 1 + k}{i} z_1^{\text{wt}v - 1 + k - i} \text{tr} Y_M(v(i)v_1, z_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \\ &= z_1^{\text{wt}v_1} \text{tr} o(v) Y_M(v_1, z_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \\ &\quad + \sum_{k \in \mathbb{Z} - \{0\}} |(\frac{z_1}{z})^k (1 - \epsilon_{v, w} \epsilon_v q^k)^{-1} \sum_{i=0}^{\infty} \sum_{m=0}^i c(\text{wt}v, i, m) k^m T(v(i)v_1) \\ &= z_1^{\text{wt}v_1} \text{tr} o(v) Y_M(v_1, z_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \\ &\quad + \sum_{i=0}^{\infty} \sum_{m=0}^i \bar{P}_{m+1}(1, \epsilon_{v, v_1} \epsilon_v, \frac{z_1}{z}, q) m! c(\text{wt}v, i, m) T(v(i)v_1) \\ &= \text{tr} o(v) o(v_1) \phi(\sigma h) q^{L(0) - c/24} + \sum_{m=0}^{\infty} \bar{P}_{m+1}(1, \epsilon_{v, v_1} \epsilon_v, \frac{z_1}{z}, q) T(v[m]v_1) \\ &= \text{tr} o(v) o(v_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} + \sum_{k=1}^{\infty} \bar{P}_k(1, \epsilon_{v, v_1} \epsilon_v, \frac{z_1}{z}, q) T(v[k-1]v_1) \end{aligned}$$

Using Lemma 6.3 and discussing the values of $\epsilon_{v, v_1} \epsilon_v$ give the result. $\square$

Sublemma 6.9 *We have*

$$T(v[-1]v_1) = \text{tr} o(v) o(v_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} + \sum_{m=1}^{\infty} E_{2m}(\tau) T(v[2m-1]v_1).$$

Proof: If $\epsilon_{v, v_1} \epsilon_v \neq 1$ then both sides of the equation are zero by Lemma 6.3. So we now assume that $\epsilon_{v, v_1} \epsilon_v = 1$. Write $v[-1]v_1 = \sum_{i \geq -1} c_i v(i)v_1$ with $c_{-1} = 1$. Then

$$\begin{aligned} T(v[-1]v_1) &= \sum_{i \geq -1} c_i z_1^{\text{wt}v + \text{wt}v_1 - i - 1} \text{tr} Y_M(v(i)v_1, z_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \\ &= \sum_{i \geq -1} c_i z_1^{\text{wt}v + \text{wt}v_1 - i - 1} \text{Res}_{z_0} z_0^i \text{tr} Y_M(Y(v, z_0)v_1, z_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \\ &= \sum_{i \geq -1} c_i z_1^{\text{wt}v + \text{wt}v_1 - i - 1} \text{Res}_{z_0} z_0^i \text{Res}_z (\frac{z_1 + z_0}{z})^{\frac{1}{2} \delta_{\bar{v}, 1}} \text{tr} X \phi(\sigma h) q^{L(0) - \frac{c}{24}} \end{aligned}$$

where

$$X = z_0^{-1} \delta\left(\frac{z - z_1}{z_0}\right) Y_M(v, z) Y_M(v_1, z_1) - \epsilon_{v, v_1} z_0^{-1} \delta\left(\frac{z_1 - z}{-z_0}\right) Y_M(v_1, z_1) Y_M(v, z).$$

Thus

$$\begin{aligned} T(v[-1]v_1) &= \sum_{i \geq -1} c_i z_1^{\text{wt}v - i - 1} \text{Res}_z \sum_{j \geq 0} \binom{\frac{1}{2}\delta_{\bar{v}, 1}}{j} z_1^{\frac{1}{2}\delta_{\bar{v}, 1} - j} z^{-\frac{1}{2}\delta_{\bar{v}, 1} - \text{wt}v} \cdot \\ &\quad \{(z - z_1)^{i+j} T(v, v_1) - \epsilon_{v, v_1} (-z_1 + z)^{i+j} T(v_1, v)\}. \end{aligned}$$

Now replace $T(v, v_1)$ and $T(v_1, v)$ with (6.14) and (6.15) respectively and note that

$$\text{tr}_M o(v_1) o(v) \phi(\sigma h) q^{L(0) - \frac{c}{24}} = \epsilon_v \text{tr}_M o(v) o(v_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}}.$$

Then $T(v[-1]v_1)$ can be written as

$$T(v[-1]v_1) = (a) + (b) \tag{6.16}$$

where

$$\begin{aligned} (a) &= \text{Res}_z z_1^{\frac{1}{2}\delta_{\bar{v}, 1} + \text{wt}v} z^{-\frac{1}{2}\delta_{\bar{v}, 1} - \text{wt}v} \{(z - z_1)^{-1} + (z_1 - z)^{-1}\} \text{tro}(v) o(v_1) \phi(\sigma h) q^{L(0) - c/24} \\ &= \text{Res}_z z_1^{\frac{1}{2}\delta_{\bar{v}, 1} + \text{wt}v} z^{-\frac{1}{2}\delta_{\bar{v}, 1} - \text{wt}v} z_1^{-1} \delta\left(\frac{z}{z_1}\right) \text{tro}(v) o(v_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \\ &= \text{tro}(v) o(v_1) \phi(\sigma h) q^{L(0) - \frac{c}{24}} \end{aligned}$$

(we have used the fact that $\frac{1}{2}\delta_{\bar{v}, 1} + \text{wt}v$ is always an integer here) and

$$\begin{aligned} (b) &= \sum_{k=1}^{\infty} \sum_{i \geq -1} c_i \text{Res}_z \sum_{j \geq 0} \binom{\frac{1}{2}\delta_{\bar{v}, 1}}{j} z_1^{\text{wt}v + \frac{1}{2}\delta_{\bar{v}, 1} - 1 - i - j} z^{-\frac{1}{2}\delta_{\bar{v}, 1} - \text{wt}v} \\ &\quad \{(z - z_1)^{i+j} P_k(1, 1, \frac{z_1}{z}, q) - (-z_1 + z)^{i+j} P_k(1, 1, \frac{z_1 q}{z}, q)\} T(v[k-1]v_1). \end{aligned}$$

From the proof of Proposition 4.3.3 of [Z] we see that

$$(b) = \sum_{k=1}^{\infty} E_{2k}(q) T(v[2k-1]v_1).$$

This completes the proof. $\square$

Lemma 6.7 now follows from Sublemma 6.9 (see the proof of Proposition 4.3.6 of [Z]).

We also need the following Lemma whose proof is contained in [Z] by using Sublemma 6.9.

Lemma 6.10 *If $v \in V$ satisfies $\sigma v = gv = hv = v$, then*

$$S(L[-2]v, \tau) = \partial S(v, \tau) + \sum_{l=2}^{\infty} E_{2l}(\tau) S(L[2l-2]v, \tau)$$

where

$$\partial S(v, \tau) = \partial_k S(v, \tau) = \frac{1}{2\pi i} \frac{d}{d\tau} S(v, \tau) + k E_2(\tau) S(L[2l-2]v, \tau)$$

Theorem 6.2 follows from Lemmas 6.3, 6.4, 6.5, 6.7 and 6.10.

Here is another important theorem whose proof is similar to that of Theorem 8.4 of [DLM3].

Theorem 6.11 *Let $M^1, M^2, \dots$ be inequivalent simple σg -twisted V -modules such that $h\sigma \circ M^i, \sigma \circ M^i$ and M^i are isomorphic. Let $T_1, T_2, \dots$ be the corresponding trace functions (6.4). Then $T_1, T_2, \dots$ are linearly independent elements of $\mathcal{C}(g, h)$.*

7 Existence of twisted modules

Although we have the definition of twisted module, but we do not know if there is an irreducible g -twisted V -module for a finite order automorphism g of V . The main result in this section gives a positive answer to the question.

Let $g, h \in \text{Aut}(V)$ commute and have finite orders.

Lemma 7.1 *Let $v \in V$ satisfy $gv = \mu^{-1}v, hv = \lambda^{-1}v$. Then the following hold:*

- (1) *The constant term of $\sum_{k=0}^{\infty} Q_k(\mu, \lambda, q)v[k-1]w$ is equal to $-v \circ_{\sigma g} w$ if $\mu \neq 1$.*
- (2) *The constant term of $\sum_{k=0}^{\infty} Q_k(\mu, \lambda, q)(L[-1]v)[k-1]w$ is equal to $-v \circ_{\sigma g} w$ if $\mu = 1, \lambda \neq 1$.*
- (3) *The constant term of $v[-2]w + \sum_{k=2}^{\infty} (2k-1)E_{2k}(q)v[2k-2]w$ is $v \circ_{\sigma g} w$ if $\mu = \lambda = 1$.*

Proof: The proof is same as that of Lemma 9.2 in [DLM3] by noting that $A_{\sigma g}(V)$ is defined by using the decomposition of V into a direct sum of eigenspaces of $\sigma(\sigma g) = g$.

□

Theorem 7.2 *Suppose that V is a simple VOSA which satisfies condition C_2 , and that $g \in \text{Aut}V$ has finite order. Then V has at least one simple g -twisted module.*

Proof: Since g is arbitrary, it is enough to prove that there exists a simple $g\sigma$ -twisted module. By Proposition 4.8, it suffices to prove that $A_{\sigma g}(V) \neq 0$.

First we show that if $\mathcal{C}(g, h) \neq 0$ for some finite order $h \in \text{Aut}(V)$ which commutes with g then $A_{g\sigma}(V)$ is nonzero. As in [DLM3] we take nonzero $S \in \mathcal{C}(g, h)$. Then there exists positive integer p such that (5.9)-(5.12) hold for all $v \in V$ with $S_p \neq 0$. We define a partial order on $\mathbb{C}$ as follows: $\lambda \geq \mu$ if $\lambda - \mu \in \frac{n}{T}$ for some nonnegative integer $n \geq 0$. We can arrange the notation so that $\lambda_{p,1}$ is minimal among all $\lambda_{p,j}$ with respect to the partial order and $a_{p,1,0}(v) \neq 0$ for some $v \in V$. Then

$$S_{p,1}(v, \tau) = \alpha(v) + \sum_{n=1}^{\infty} a_{p,1,n}(v)q^{n/T} \quad (7.1)$$

defines a function $\alpha : V \rightarrow \mathbb{C}$ which is not identically zero. Because $S(v, \tau)$ is linear in v , α is a linear function on V . Furthermore, α vanishes on $O_{\sigma g}(V)$ (see the proof of Lemma 9.3 of [DLM3]). Thus $A_{g\sigma}(V)$ is nonzero.

Since V is a simple V -module by the assumption, $T(v) = \text{tr}_V o(v) g \sigma q^{L(0)-c/24}$ defines a nonzero element in $\mathcal{C}(\sigma, g)$ by Theorems 6.2 and 6.11. Since $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ induces a linear isomorphism between $\mathcal{C}(\sigma, g)$ and $\mathcal{C}(g, \sigma)$ by Theorem 5.4. In particular, $\mathcal{C}(g, \sigma)$ is nonzero. The proof is complete. $\square$

8 The main theorems

Recall Theorem 6.11. In the case V is $g\sigma$ -rational, we can strength the results obtained in 6.11.

Theorem 8.1 *Suppose that V is σg -rational and C_2 -cofinite. Let $M^1, \dots, M^m$ be all of the inequivalent, simple σg -twisted V -modules such that $h \circ M^i$, $\sigma \circ M^i$ and M^i are isomorphic. Let $T_1, \dots, T_m$ be the corresponding trace functions defined by (6.4). Then $T_1, \dots, T_m$ form a basis of $\mathcal{C}(g, h)$.*

Remark 8.2 *In the proof of Theorem 8.1, we only use a consequence of σg -rationality of $V - A_{\sigma g}(V)$ is finite dimensional semisimple algebra instead of σg -rationality of V itself. So we can replace the σg -rationality assumption by the semi-simplicity of $A_{\sigma g}(V)$.*

We begin with an arbitrary function $S \in \mathcal{C}(g, h)$. We have already seen that S can be represented as

$$S(v, \tau) = \sum_{i=0}^p (\log q_{\frac{1}{T}})^i S_i(v, \tau) \quad (8.1)$$

for fixed p and all $v \in V$, with each S_i satisfying (5.10)-(5.12). Then Theorem 8.1 follows from the following propositions

Proposition 8.3 (1) *Each S_i is a linear combination of the functions $T_1, \dots, T_m$.*
(2) *$S_i = 0$ if $i > 0$.*

Using the proof of Proposition 10.5 in [DLM3] we can show that Proposition (2) follows from (1). So we only need to prove (1).

We assume the setting in Section 7. In particular, $S_p \neq 0$, each $S_{p,j} \neq 0$ (5.10), and $\alpha : V \rightarrow \mathbb{C}$ vanishes on $O_g(V)$ which induces a linear function

$$\alpha : A_{g\sigma}(V) \rightarrow \mathbb{C}.$$

Lemma 8.4 *Suppose that $u, v \in V$ and satisfy $hu = \rho u$, $hv = \lambda v$, $\rho, \lambda \in \mathbb{C}$. Then*

$$\alpha(u *_{g\sigma} v) = \epsilon_{u,v} \rho \delta_{\rho\sigma, 1} \alpha(v *_{g\sigma} u). \quad (8.2)$$

Proof: Let $gu = \xi u$ and $gv = \nu v$ for scalars ξ, ν . If ξ or ν is not equal to 1 then u (resp. v) lies in $O_{g\sigma}(V)$ by Lemma 3.1 of [DZ]. This implies that $u *_{g\sigma} v$ and $v *_{g\sigma} u$ are zero by Theorem 3.3 of [DZ].

We now assume $gu = u, gv = v$. It is clear that $u *_g v$ is an eigenvector for h with eigenvalue $\rho\lambda$. If $\rho\lambda \neq 1$ then $u *_g v$ and $v *_g u$ lie in $O(g, h)$ by (5.3). Then $S(u *_g v) = S(v *_g u) = 0$ by (C3). This leads to both sides of (8.2) being 0. So finally we can assume that $\rho\lambda = 1$.

Recall the decomposition (4.1). Then $u, v \in V^0$. By Lemma 3.3 of [DZ],

$$u *_g v - (-1)^{\tilde{u}\tilde{v}} v *_g u \equiv \text{Res}_z Y(u, z) v (1+z)^{\text{wt}u-1} \pmod{O(V^0)}.$$

By (3.11)

$$u *_g v - (-1)^{\tilde{u}\tilde{v}} v *_g u \equiv \sum_{i=0}^{\infty} \binom{\text{wt}u-1}{i} u(i)v \equiv u[0]v \pmod{O(V^0)}. \quad (8.3)$$

Since $O(V^0) \subset O_{g\sigma}(V)$, $u[0]v \in O(g, h)$ if $\rho = 1$ by (5.1). Thus $\alpha(u *_g v - \epsilon_{u,v} v *_g u) = 0$ if $\rho = 1$.

We now deal with the case $\rho \neq 1$. From the proof of Lemma 9.2 of [DLM3], we see that the constant term of

$$\sum_{k=0}^{\infty} Q_k(1, \rho^{-1}, q) u[k-1]v$$

is

$$-u *_g v + \frac{1}{1-\rho^{-1}} u[0]v.$$

The vanishing condition of S on $\sum_{k=0}^{\infty} Q_k(1, \rho^{-1}, q) u[k-1]v \in O(g, h)$ yields

$$\alpha(u *_g v) = \frac{1}{1-\rho^{-1}} \alpha(u[0]v).$$

Using (8.3) gives the desired result. $\square$

Lemma 8.5 *Suppose that $u, v \in V$ with $h\sigma u = \rho'u, h\sigma v = \lambda'v, \rho', \lambda' \in \mathbb{C}$. Then*

$$\alpha(u *_g v) = \rho' \delta_{\rho'\lambda', 1} \alpha(v *_g u). \quad (8.4)$$

Proof: We still assume that $hu = \rho u, hv = \lambda v$. Then $\rho' = \epsilon_u \rho$ and $\lambda' = \epsilon_v \lambda$. By Lemma 8.4 we have

$$\alpha(u *_g v) = \epsilon_{u,v} \epsilon_u \rho' \delta_{\epsilon_u \epsilon_v \rho' \lambda', 1} \alpha(v *_g u).$$

If $\epsilon_u \epsilon_v = -1$, then $\sigma(u *_g v) = -u *_g v$. By (5.3), both $u *_g v$ and $v *_g u$ lie in $O(g, h)$. This implies that $S(u *_g v) = S(v *_g u) = 0$. As a result, both sides of (8.4) are equal to 0. If $\epsilon_u \epsilon_v = 1$, then $\epsilon_{u,v} \epsilon_u = 1$ and (8.4) still holds. $\square$

The rest of the proof of Proposition 8.3 (1) follows from the argument given in Section 10 of [DLM3].

As in [DLM3] we have important corollaries.

Theorem 8.6 *Suppose that V is rational and C_2 -cofinite.*

(1) *If the group $\langle g, h \rangle$ generated by g and h is cyclic with generator k . Then the dimension of $\mathcal{C}(g, h)$ is equal to the number of inequivalent, k, σ -stable, σ -twisted simple V -modules.*

(2) *The number of inequivalent, σ -stable simple g -twisted V -modules is at most equal to the number of g, σ -stable, simple V -modules, with equality if V is g -rational.*

The proof is the same as that of Theorem 10.2 in [DLM3] by using Lemma 6.1 and Theorem 8.1.

As in the theory of vertex operator algebra, a simple vertex operator superalgebra V is called *holomorphic* in case V is rational and if V is the unique simple V -module.

Theorem 8.7 *Suppose that V is holomorphic and is C_2 -cofinite. For each automorphism g of V of finite order, there is a unique σ -stable simple g -twisted V -module $V(g)$. Moreover if $\langle g, h \rangle$ is cyclic then $\mathcal{C}(g, h)$ is spanned by $T_{V(\sigma g)}(v, g, h, q)$.*

It is proved in [Z] that if a vertex operator algebra V is rational and C_2 -cofinite, then the space spanned by $T_M(v, q) = \text{tr}_M o(v) q^{L(0)-c/24}$ with M running through the inequivalent simple V -modules admits a representation of the modular group Γ . But this is not true anymore in the present situation. In fact, if V is holomorphic such that $V_1 \neq 0$, then the character $\text{tr}_V q^{L(0)-c/24}$ of V is not an eigenvector for the matrix $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. Here is the corresponding result for the vertex operator superalgebra, which is an immediate consequence of Theorems 6.2 and 8.1.

Theorem 8.8 *Let V be σ -rational and C_2 -cofinite VOSA. Let $M^1, \dots, M^m$ be all of the inequivalent, simple σ -twisted V -modules such that $\sigma \circ M^i$ and M^i are isomorphic. Then the space spanned by*

$$T_i(v, \tau) = T_i(v, (1, 1), \tau) = \text{tr}_{M^i} o(v) \phi(\sigma) q^{L(0)-c/24}$$

admits a representation of the modular group. That is, for any $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ there exists a $m \times m$ invertible complex matrices (γ_{ij}) such that

$$T_i(v, \frac{a\tau + b}{c\tau + d}) = (c\tau + d)^n \sum_{j=1}^m \gamma_{ij} T_j(v, \tau)$$

for all $v \in V_{[n]}$. Moreover, the matrix (γ_{ij}) is independent of v .

Finally we discuss the rationality of the central charge c and the conformal weights of a g -twisted module under certain conditions. Let M be a simple g -twisted V -module. Recall from (4.8) that M is direct sum of $L(0)$ -eigenspaces

$$M = \bigoplus_{n=0}^{\infty} M_{\lambda+n/T'}$$

for some $\lambda \in \mathbb{C}$ such that $M_{\lambda} \neq 0$. The λ is called the conformal weight of M .

Theorem 8.9 *Assume that V is C_2 -cofinite and $g \in \text{Aut}V$ has finite order. Suppose that V is $\sigma^i g^j$ -rational for all integers i, j . Then each σ -stable simple $\sigma^i g^j$ -twisted V -module has rational conformal weight, and the central charge c of V is rational.*

The proof of this theorem is the same as that of Theorem 11.2 in [DLM3] (also see [AM]) using Lemma 6.1.

9 Modular invariance over Γ_θ

In order to get the modular invariance over the full modular group Γ we only considered the σ -stable twisted modules in previous sections. In this section we remove the σ -stable assumption. But we have a modular invariance over the Γ_θ instead of the full modular group Γ as in [H]. Since the proofs of most results in this section are very similar to those given in the previous sections and [DLM3] we present the results without proofs.

Recall that G is a finite automorphism group of V . Let $g \in G$ and M a simple g -twisted V -module. Also recall $M \circ h$ for $h \in G$. Let $G_M = \{h \in G | M \circ h \cong M\}$ be the stablizer of M . Then M is a projective module for G_M , denoting by ϕ the projective action. For $h \in G_M$ and $v \in V$ set

$$F_M(v, (g, h), q) = \text{tr}_M o(v) \phi(h) q^{L(0)-c/24}.$$

Theorem 9.1 *Let V be a rational and C_2 -cofinite vertex operator superalgebra.*

(1) *If g is an automorphism of V of finite order then the number of inequivalent, g -stable, simple g -twisted V -modules is at most equal to the number of g -stable simple untwisted V -modules. If V is g -rational, the number of inequivalent, g -stable simple g -twisted V -modules is precisely the number of g -stable simple untwisted V -modules.*

(2) *Let $g \in \text{Aut}V$ have finite order. Suppose that V is g^i -rational for all integers i . Then each g -stable simple g^i -twisted V -module has rational conformal weight, and the central charge c of V is rational.*

(3) *Let G be a finite automorphism group of V . Let $g \in G$ and M a simple g -twisted V -module and $h \in G_M$. Then the trace function $F_M(v, (g, h), q)$ converges to a holomorphic function in the upper half plane $\mathfrak{h}$.*

(4) *Assume that V is x -rational for each $x \in G$. Let $v \in V$ satisfy $\text{wt}[v] = k$. Then the space of (holomorphic) functions in $\mathfrak{h}$ spanned by the trace functions $F_M(v, (g, h), \tau)$ for all choices of g, h in G and h -stable M is a (finite-dimensional) Γ_θ -module. That is, if $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_\theta$ then we have an equality*

$$F_M(v, (g, h), \frac{a\tau + b}{c\tau + d}) = (c\tau + d)^k \sum_W \gamma_{M,W} F_W(v, (g^a h^c, g^b h^d), \tau),$$

The statements (3) and (4) are exactly the same as in [DLM3] for vertex operator algebras except that we use the group Γ_θ here instead of Γ in [DLM3]. In the case $G = 1$, Parts (3) and (4) have been obtained previously in [H].

10 Examples

In this section we will use the examples in [DZ] (cf. Theorem 7.1, propositions 7.2 and 7.4) to show the modular invariance. So we are working in the setting of Section 7 of [DZ]. In particular, H is a vector space of dimension l which we assume to be even, $V(H, \mathbb{Z} + \frac{1}{2})$ is a VOSA with central charge $c = \frac{l}{2}$ and $V(H, \mathbb{Z})$ is the unique irreducible σ -twisted module. We are going to study the trace functions $T(\mathbf{1}, (\sigma^i, \sigma^j), q)$ which is the graded σ^{j+1} -trace on the irreducible σ^{i+1} -twisted module for $V(H, \mathbb{Z} + \frac{1}{2})$ for $i, j = 0, 1$.

We first recall the important modular form

$$\eta(\tau) = q^{\frac{1}{24}} \prod_{n \geq 1} (1 - q^n).$$

The trace functions in our examples can be expressed as rational functions in $\eta(\tau)$. Let $S = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ and $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ be the standard generators of $SL(2, \mathbb{Z})$. Then

$$\eta\left(\frac{-1}{\tau}\right) = (-i\tau)^{\frac{1}{2}} \eta(\tau)$$

and

$$\eta(\tau + 1) = e^{\frac{\pi i}{12}} \eta(\tau).$$

Note that the conformal weight of $V(H, \mathbb{Z})$ is $\lambda = \frac{l}{16}$. It is easy to compute that

$$\begin{aligned} T(\mathbf{1}, (1, 1), \tau) &= \text{tr}_{V(H, \mathbb{Z})} \sigma q^{L(0) - \frac{l}{48}} = q^{\frac{l}{24}} \prod_{m=1}^{l/2} (1 + (-1)^m) \prod_{n>0} (1 - q^n)^l = 0 \\ T(\mathbf{1}, (1, \sigma), \tau) &= \text{tr}_{V(H, \mathbb{Z})} q^{L(0) - \frac{l}{48}} = 2^{l/2} q^{\frac{l}{24}} \prod_{n=1}^{\infty} (1 + q^n)^l = 2^{l/2} \left[\frac{\eta(2\tau)}{\eta(\tau)} \right]^l \\ T(\mathbf{1}, (\sigma, 1), \tau) &= \text{tr}_{V(H, \frac{1}{2} + \mathbb{Z})} \sigma q^{L(0) - \frac{l}{48}} = q^{-\frac{l}{48}} \prod_{n=0}^{\infty} (1 - q^{n+\frac{1}{2}})^l = \left[\frac{\eta(\tau/2)}{\eta(\tau)} \right]^l \\ T(\mathbf{1}, (\sigma, \sigma), \tau) &= \text{tr}_{V(H, \frac{1}{2} + \mathbb{Z})} q^{L(0) - \frac{l}{48}} = q^{-\frac{l}{48}} \prod_{n=0}^{\infty} (1 + q^{n+\frac{1}{2}})^l = \left[\frac{\eta(\tau)^2}{\eta(\tau/2)\eta(2\tau)} \right]^l. \end{aligned}$$

The modular transformation law then gives the modular transformation property of functions $T(\mathbf{1}, (\sigma^i, \sigma^j), \tau)$ for $i, j = 0, 1$:

$$T(\mathbf{1}, (\sigma^i, \sigma^j), \frac{a\tau + b}{c\tau + d}) = \gamma_{i,j} T(\mathbf{1}, (\sigma^{ai+cj}, \sigma^{bi+dj}), \tau)$$

for some constant $\gamma_{i,j} \in \mathbb{C}$ where $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$. The result matches what Theorems 5.4 and 8.1 assert.

Here is another example. We still take V to be $V(H, \frac{1}{2} + \mathbb{Z})$ with $l = \dim H$ being even. Let g be an isometry of H such that $g(h_1) = -h_2, g(h_2) = -h_1, g(h_1^*) = -h_2^*, g(h_2^*) = -h_1^*$.

And $g(h_i) = -h_i, g(h_i^*) = -h_i^*$ for $i = 3, \dots, l/2$ where $\{h_1, \dots, h_{l/2}, h_1^*, \dots, h_{l/2}^*\}$ is a basis of H such that $(h_i, h_j) = (h_i^*, h_j^*) = 0$ and $(h_i, h_j^*) = \delta_{i,j}$. Then $g\sigma$ interchanges h_1 and h_2 , h_1^* and h_2^* , and leaves other h_i, h_i^* invariant. Then we have

$$H = H^{0*} \oplus H^{1*}, \quad H = H^0 \oplus H^1$$

where $g\sigma|_{H^{i*}} = (-1)^i$ and $g|_{H^i} = (-1)^i$. It is easy to see that

$$H^{0*} = \mathbb{C}(h_1 + h_2) \oplus \mathbb{C}(h_1^* + h_2^*) \oplus \bigoplus_{j=3}^{l-2} (\mathbb{C}h_j + \mathbb{C}h_j^*),$$

$$H^{1*} = \mathbb{C}(h_1 - h_2) \oplus \mathbb{C}(h_1^* - h_2^*),$$

and $H^0 = H^{1*}, H^1 = H^{0*}$. Note that g is lifted to an automorphism of the vertex operator superalgebra $V(H, \mathbb{Z} + \frac{1}{2})$ in an obvious way.

By proposition 7.4 in [DZ],

$$M = \wedge[(h_1 - h_2)(-n), (h_1^* - h_2^*)(-m), h(-m - \frac{1}{2}) | n, m \in \mathbb{Z}, n > 0, m \geq 0, h \in H^1]$$

is the unique irreducible $g\sigma$ -twisted $V(H, \mathbb{Z} + \frac{1}{2})$ -module whose conformal weight is $\frac{1}{8}$. Thus

$$\begin{aligned} T(\mathbf{1}, (g, \sigma), \tau) &= \text{tr}_M q^{L(0) - \frac{l}{48}} \\ &= \sum_{n \in \frac{1}{2}\mathbb{Z}} (\dim M_{n+\frac{1}{8}}) q^{n - \frac{l}{48}} \\ &= 2 \left[\prod_{0 < n \in \mathbb{Z}} (1 + q^n) \right]^2 \left[\prod_{0 \leq n \in \mathbb{Z}} (1 + q^{n+\frac{1}{2}}) \right]^{l-2} q^{\frac{l}{8} - \frac{l}{48}} \\ &= 2 \left[q^{-\frac{1}{24}} \frac{\eta(2\tau)}{\eta(\tau)} \right]^2 \left[q^{\frac{1}{48}} \frac{\eta^2(\tau)}{\eta(\frac{\tau}{2})\eta(2\tau)} \right]^{l-2} q^{\frac{l}{8} - \frac{l}{48}} \\ &= 2 \frac{\eta(\tau)^{2l-6}}{\eta(2\tau)^{l-4} \eta(\frac{\tau}{2})^{l-2}}, \end{aligned}$$

$$\begin{aligned} T(\mathbf{1}, (\sigma, g), \tau) &= \text{tr}_{V(H, \mathbb{Z} + \frac{1}{2})} g\sigma q^{L(0) - \frac{l}{48}} \\ &= \sum_{0 \leq n \in \frac{1}{2}\mathbb{Z}} (\text{tr}_{V(H, \mathbb{Z} + \frac{1}{2})_n} g\sigma) q^{n - \frac{l}{48}} \\ &= \left[\prod_{0 \leq n \in \mathbb{Z}} (1 - q^{n+\frac{1}{2}}) \right]^2 \left[\prod_{0 \leq n \in \mathbb{Z}} (1 + q^{n+\frac{1}{2}}) \right]^{l-2} q^{-\frac{l}{48}} \\ &= \left[q^{\frac{1}{48}} \frac{\eta(\frac{\tau}{2})}{\eta(\tau)} \right]^2 \left[q^{\frac{l}{48}} \frac{\eta^2(\tau)}{\eta(\frac{\tau}{2})\eta(2\tau)} \right]^{l-2} q^{-l/48} \\ &= \frac{\eta(\tau)^{2l-6}}{\eta(\frac{\tau}{2})^{l-4} \eta(2\tau)^{l-2}}. \end{aligned}$$

and

$$\begin{aligned}
T(\mathbf{1}, (g, g\sigma), \tau) &= \text{tr}_M g q^{L(0) - \frac{l}{48}} \\
&= \sum_{0 \leq n \in \frac{1}{2}\mathbb{Z}} (\text{tr}_{M_{n+\frac{1}{8}}} g) q^{L(0) - \frac{l}{48}} \\
&= 2q^{\frac{1}{8} - \frac{l}{48}} \left[\prod_{0 < n \in \mathbb{Z}} (1 + q^n) \right]^2 \left[\prod_{0 \leq n \in \mathbb{Z}} (1 - q^{n+\frac{1}{2}}) \right]^{l-2} \\
&= 2q^{\frac{1}{8} - \frac{l}{48}} \left[q^{\frac{-1}{24}} \frac{\eta(2\tau)}{\eta(\tau)} \right]^2 \left[q^{\frac{1}{48}} \frac{\eta(\frac{\tau}{2})}{\eta(\tau)} \right]^{l-2} \\
&= 2 \frac{\eta(2\tau)^2 \eta(\frac{\tau}{2})^{l-2}}{\eta(\tau)^l}
\end{aligned}$$

Then

$$\begin{aligned}
T(\mathbf{1}, (g, \sigma), S\tau) &= 2 \frac{\eta(-\frac{1}{\tau})^{2l-6}}{\eta(-\frac{2}{\tau})^{l-4} \eta(\frac{-1}{2\tau})^{l-2}} \\
&= 2 \frac{(-i\tau)^{\frac{2l-6}{2}} \eta(\tau)^{2l-6}}{(-i\tau/2)^{\frac{l-4}{2}} \eta(\frac{\tau}{2})^{l-4} (-i2\tau)^{\frac{l-2}{2}} \eta(2\tau)^{l-2}} \\
&= 2\mu \frac{\eta(\tau)^{2l-6}}{\eta(\frac{\tau}{2})^{l-4} \eta(2\tau)^{l-2}}
\end{aligned}$$

where μ is a root of unity. This implies that

$$T(\mathbf{1}, (g, \sigma), S\tau) = 2\mu T(\mathbf{1}, (g, \sigma)S, \tau) = 2\mu T(\mathbf{1}, (\sigma, g), \tau).$$

Using the fact that

$$\eta\left(\frac{\tau+1}{2}\right) = \eta(\tau) \frac{\eta(\tau)^2}{\eta(\frac{\tau}{2})\eta(2\tau)}$$

we have

$$\begin{aligned}
T(\mathbf{1}, (g, \sigma), T\tau) &= T(\mathbf{1}, (g, \sigma), \tau + 1) \\
&= 2 \frac{\eta(\tau + 1)^{2l-6}}{\eta(2(\tau + 1))^{l-4} \eta(\frac{\tau+1}{2})^{l-2}} \\
&= 2\nu \frac{\eta(\tau)^{2l-6}}{\eta(2\tau)^{l-4}} \frac{\eta(\frac{\tau}{2})^{l-2} \eta(2\tau)^{l-2}}{\eta(\tau)^{2l-4}} \\
&= 2\nu \frac{\eta(2\tau)^2 \eta(\frac{\tau}{2})^{l-2}}{\eta(\tau)^l}
\end{aligned}$$

for some root of unity ν . Since $(g, \sigma)T = (g, g\sigma)$, we see that

$$T(\mathbf{1}, (g, \sigma), T\tau) = \nu T(\mathbf{1}, (g, g\sigma), \tau).$$

This again verifies Theorems 5.4 and 8.1 in this special case. The reader could determine $T(v, (h, k), \tau)$ for other vector $v \in V$ in these two cases, but the computation will be more complicated and difficult.

References

- [ABD] T. Abe, G. Buhl and C. Dong, Rationality, regularity and C_2 -cofiniteness, *Trans. AMS.* **356** (2004), 3391-3402.
- [AM] G. Anderson and G. Moore, Rationality in conformal field theory, *Comm. Math. Phys.* **117** (1988), 441-450.
- [B1] R. E. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, *Proc. Natl. Acad. Sci. USA* **83** (1986), 3068-3071.
- [B2] R. E. Borcherds, Monstrous moonshine and monstrous Lie superalgebras, *Invent. Math.* **109** (1992), 405-444.
- [CN] J. H. Conway and S. P. Norton, Monstrous Moonshine, *Bull. London. Math. Soc.* **12** (1979), 308-339.
- [DVVV] R. Dijkgraaf, C. Vafa, E. Verlinde and H. Verlinde, The operator algebra of orbifold models, *Comm. Math. Phys.* **123** (1989), 485-526.
- [D] C. Dong, Twisted modules for vertex algebras associated with even lattice, *J. of Algebra* **165** (1994), 91-112.
- [DL] C. Dong and J. Lepowsky, Generalized Vertex Algebras and Relative Vertex Operators, *Progress in Math.* Vol. 112, Birkhäuser, Boston, 1993.
- [DLM1] C. Dong, H. Li and G. Mason, Regularity of rational vertex operator algebras, *Advances. in Math.* **132** (1997), 148-166.
- [DLM2] C. Dong, H. Li and G. Mason, Twisted representations of vertex operator algebras, *Math. Ann.* **310** (1998), 571-600.
- [DLM3] C. Dong, H. Li and G. Mason, Modular invariance of trace functions in orbifold theory and generalized moonshine, *Comm. Math. Phys.* **214** (2000), 1-56.
- [DM1] C. Dong and G. Mason, Rational vertex operator algebras and the effective central charge, *International Math. Research Notices* **56** (2004), 2989-3008.
- [DM2] C. Dong and G. Mason, Holomorphic vertex operator algebras of small central charges, *Pacific J. Math.* **213** (2004), 253-266.
- [DZ] C. Dong and Z. Zhao, Twisted representations of vertex operator superalgebras, preprint.
- [FFR] A. J. Feingold, I. B. Frenkel and J. F. X. Ries, Spinor construction of vertex operator algebras, triality and $E_8^{(1)}$, *Contemporary Math.* **121**, 1991.

- [FLM1] I. B. Frenkel, J. Lepowsky and A. Meurman, A natural representation of the Fischer-Griess Monster with the modular function J as character, *Proc. Natl. Acad. Sci. USA* **81** (1984), 3256-3260.
- [FLM2] I. B. Frenkel, J. Lepowsky and A. Meurman, Vertex operator calculus, in: *Mathematical Aspects of String Theory, Proc. 1986 Conference, San Diego*. ed. by S.-T. Yau, World Scientific, Singapore, 1987, 150-188.
- [FLM3] I. B. Frenkel, J. Lepowsky and A. Meurman, Vertex Operator Algebras and the Monster, *Pure and Applied Math.*, Vol. **134**, Academic Press, 1988.
- [G] R. Griess, The Friendly Giant, *Invent. Math.* **69** (1982), 1-102.
- [H] G. Höhn, Self-dual vertex operator superalgebras and the Baby Monster, Bonn Mathematical Publications, 286. Universitt Bonn, Mathematisches Institut, Bonn, 1996.
- [Hu] Y. Huang, Vertex operator algebras and the Verlinde conjecture, math.QA/0406291.
- [L] J. Lepowsky, Calculus of twisted vertex operators, *Proc. Natl. Acad. Sci. USA* **82** (1985), 8295-8299.
- [Li] H. Li, Local systems of vertex operators, vertex superalgebras and modules, *J. Pure Appl. Alg.* **109** (1996) 143-195.
- [M1] M. Miyamoto, A modular invariance on the theta functions defined on vertex operator algebras, *Duke Math. J.* **101** (2000), 221-236.
- [M2] M. Miyamoto, Modular invariance of vertex operator algebras satisfying C_2 -cofiniteness *Duke Math. J.* **122** (2004), 51-91.
- [Mo] P. Montague, Orbifold constructions and the classification of self-dual $c = 24$ conformal field theory, *Nucl. Phys.* **B428** (1994), 233-258.
- [MS] G. Moore and N. Seiberg, Classical and quantum conformal field theory, *Comm. Math. Phys.* **123** (1989), 177-254.
- [N] S. Norton, Generalized moonshine, *Proc. Symp. Pure. Math., American Math. Soc.* **47** (1987), 208-209.
- [S] A. N. Schellekens, Meromorphic $c = 24$ conformal field theories, *Comm. Math. Phys.* **153** (1993), 159.
- [T1] M. Tuite, Monstrous moonshine from orbifolds, *Comm. Math. Phys.* **146** (1992), 277-309.

- [T2] M. Tuite, On the relationship between monstrous moonshine and the uniqueness of the moonshine module, *Comm. Math. Phys.* **166** (1995), 495-532.
- [T3] M. Tuite, Generalized moonshine and abelian orbifold constructions, *Contemp. Math.* **193** (1996), 353-368.
- [V] E. Verlinde, Fusion rules and modular transformations in 2D conformal field theory *Nucl. Phys.* **B300** (1980), 360-376.
- [Y] H. Yamauchi, Modularity on vertex operator algebras arising from semisimple primary vectors, *Internat. J. Math.* **15** (2004), 87-109.
- [Z] Y. Zhu, Modular invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.* **9** (1996), 237-302.